



## Research Article

# Survival of White-Tailed Deer Neonates in Minnesota and South Dakota

TROY W. GROVENBURG,<sup>1</sup> *Department of Wildlife and Fisheries Sciences, South Dakota State University, Brookings, SD 57007, USA*

CHRISTOPHER C. SWANSON, *Department of Wildlife and Fisheries Sciences, South Dakota State University, Brookings, SD 57007, USA*

CHRISTOPHER N. JACQUES, *Bureau of Science Services, Wisconsin Department of Natural Resources, 2801 Progress Road, Madison, WI 53716, USA*

ROBERT W. KLAVER, *US Geological Survey Center for Earth Resources Observation and Science (EROS), 47914 252nd Street, Sioux Falls, SD 57198, USA*

TODD J. BRINKMAN, *Department of Biology and Wildlife, Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK 99775, USA*

BENJAMIN M. BURRIS, *Department of Wildlife and Fisheries Sciences, South Dakota State University, Brookings, SD 57007, USA*

CHRISTOPHER S. DEPERNO, *Department of Forestry and Environmental Resources, Fisheries and Wildlife Sciences Program, North Carolina State University, Raleigh, NC 27695, USA*

JONATHAN A. JENKS, *Department of Wildlife and Fisheries Sciences, South Dakota State University, Brookings, SD 57007, USA*

**ABSTRACT** Understanding the influence of intrinsic (e.g., age, birth mass, and sex) and habitat factors on survival of neonate white-tailed deer improves understanding of population ecology. During 2002–2004, we captured and radiocollared 78 neonates in eastern South Dakota and southwestern Minnesota, of which 16 died before 1 September. Predation accounted for 80% of mortality; the remaining 20% was attributed to starvation. Canids (coyotes [*Canis latrans*], domestic dogs) accounted for 100% of predation on neonates. We used known fate analysis in Program MARK to estimate survival rates and investigate the influence of intrinsic and habitat variables on survival. We developed 2 a priori model sets, including intrinsic variables (model set 1) and habitat variables (model set 2; forested cover, wetlands, grasslands, and croplands). For model set 1, model { $S_{\text{age-interval}}$ } had the lowest AIC<sub>c</sub> (Akaike's information criterion for small sample size) value, indicating that age at mortality (3-stage age-interval: 0–2 weeks, 2–8 weeks, and >8 weeks) best explained survival. Model set 2 indicated that habitat variables did not further influence survival in the study area;  $\beta$ -estimates and 95% confidence intervals for habitat variables in competing models encompassed zero; thus, we excluded these models from consideration. Overall survival rate using model { $S_{\text{age-interval}}$ } was 0.87 (95% CI = 0.83–0.91); 61% of mortalities occurred at 0–2 weeks of age, 26% at 2–8 weeks of age, and 13% at >8 weeks of age. Our results indicate that variables influencing survival may be area specific. Region-specific data are needed to determine influences of intrinsic and habitat variables on neonate survival before wildlife managers can determine which habitat management activities influence neonate populations. © 2011 The Wildlife Society.

**KEY WORDS** habitat, landscape, Minnesota, mortality, neonate, *Odocoileus virginianus*, predation, South Dakota, survival.

Understanding white-tailed deer (*Odocoileus virginianus*) population dynamics requires knowledge of survival rates and cause-specific mortality (Nelson and Mech 1986, Dusek et al. 1992, DePerno et al. 2000, DelGiudice et al. 2002). Research on neonates (<1 month of age) provides information regarding reproduction, sex ratios, mortality, movements, and behavior (Downing and McGinnes 1969). Furthermore, knowledge of neonate mortality is critical to understanding how pre-hunting season survival rates affect deer harvest strategies (Porath 1980). However, neonates rely on cryptic coloration and inactivity making capture difficult and survival information costly to collect (Porath 1980). Therefore, biologists often make educated guesses pertaining to neonate survival. Nevertheless, knowledge of female behavior (Downing and McGinnes 1969, Huegel et al. 1985a) and recent technology (e.g., vaginal-implant transmitters [VITs]; Bowman and Jacobson 1998, Carstensen et al. 2003, Swanson et al. 2008) improve capture success of neonate white-tailed deer.

Previous research indicates survival rates and cause-specific mortality of deer differ regionally and seasonally with respect to sex, age-class, and density of deer (Gavin et al. 1984, Dusek et al. 1992, Whitlaw et al. 1998, DelGiudice et al. 2002). Sources of mortality include starvation (Carroll and Brown 1977), disease (Cook et al. 1971, Schulz et al. 1983, Brinkman et al. 2004a), and predation (Huegel et al. 1985a, Nelson and Woolf 1987, Kunkel and Mech 1994, Rohm et al. 2007).

Macrohabitat variables affect survival of other wildlife species (Brown and Litvaitis 1995, Stuart-Smith et al. 1997, Kunkel and Pletscher 2000, Thogmartin and Schaeffer 2000) and clearly influence neonate survival by affecting predator distribution, density, and hunting efficiency (Gese et al. 1996, Dijak and Thompson 2000, Rohm et al. 2007). In Illinois, Rohm et al. (2007) documented that intrinsic and macrohabitat variables influenced survival, whereas in Pennsylvania, Vreeland et al. (2004) provided contradictory results relative to these variables. However, Vreeland et al. (2004) only evaluated 4 variables (i.e., habitat edge density, habitat diversity, proportion of herbaceous habitat, and road density), whereas Rohm et al. (2007) considered >10 covariates.

Limited survival information exists for neonate white-tailed deer in the Prairie Coteau of eastern South Dakota

Received: 21 December 2009; Accepted: 4 May 2010


<sup>1</sup>E-mail: troy.grovenburg@sdsu.edu

and the Minnesota River Valley of southwest Minnesota. Brinkman et al. (2004a) documented survival and cause-specific mortality for neonates in Lincoln and Pipestone counties in southwestern Minnesota but did not investigate the influence of intrinsic and macrohabitat variables on survival. Our objectives were to 1) estimate survival and document cause-specific mortality of neonate white-tailed deer in eastern South Dakota and southwest Minnesota and 2) determine the influence of intrinsic and habitat characteristics on neonate survival in an intensively farmed landscape.

## STUDY AREA

We conducted our study in eastern South Dakota and southwest Minnesota (Fig. 1). The region was fragmented and dominated by intense row-crop agriculture. Study sites we selected for neonate captures included Brookings County, South Dakota and Lincoln, Pipestone, Redwood, and

Renville counties, Minnesota (Fig. 1). Brookings, Pipestone, and Lincoln counties occurred within the Prairie Coteau physiographic region, whereas, Redwood and Renville counties occurred within the Minnesota River Valley. As a result of Wisconsin Glaciation (10,000–100,000 yr ago), lobes of glacial ice shaped the landscape in this region (Minnesota River Basin Data Center 2003), creating steep topography along the Prairie Coteau and Minnesota River and its tributaries (Voigtlander 1999) compared to upland prairie habitats. The Minnesota River Valley was a linear corridor heavily forested with small interspersed grassland remnants and adjacent lands comprised primarily of cultivated crops. The Coteau was a large, dead-ice moraine that historically contained numerous wetlands (Johnson and Larson 1999). In eastern South Dakota, approximately 35% of natural wetlands were drained through anthropogenic modifications (e.g., agriculture; Dahl 1990, Johnson and Higgins 1997).



**Figure 1.** Neonate white-tailed deer (*Odocoileus virginianus*) study area (shaded) in eastern South Dakota and southwestern Minnesota, USA, 2001–2004.

Land-cover types in this region were primarily agricultural, with cultivated land and pasture–grassland dominating the landscape. Land cover of the study sites on the Prairie Coteau was 58.9% cultivated, 31.4% pasture–grassland, 4.2% wetland, and 2.3% forest (Smith et al. 2002). The Minnesota River Valley study sites were composed of 85.8% cultivated, 7.0% pasture–grassland, 3.0% forest, and 2.4% wetland (Vogelmann et al. 2001, Smith et al. 2002). Harvested crops included soybeans, corn, wheat, oats, and alfalfa. Grasslands in this region were dominated by tall and mixed-prairie grasses, including Indiangrass (*Sorghastrum nutans*), switchgrass (*Panicum virgatum*), big bluestem (*Andropogon gerardii*), little bluestem (*Schizachyrium scoparium*), smooth bromegrass (*Bromus inermis*), tall dropseed (*Sporobolus asper*), prairie dropseed (*S. heterolepis*), sideoats grama (*Bouteloua curtipendula*), porcupine grass (*Stipa spartea*), and western wheatgrass (*Elymus smithii*; Johnson and Larson 1999). In low, wet areas, prairie cordgrass (*Spartina pectinata*), reedgrass (*Calamagrostis arundinacea*), reed canarygrass (*Phalaris arundinacea*), common reed (*Phragmites australis*), cattails (*Typha* spp.), rushes (*Juncus* spp.), and sedges (Cyperaceae) were common species on the landscape (Johnson and Larson 1999, Voigtlander 1999). Woodlands were dominated by American elm (*Ulmus americana*), bur oak (*Quercus macrocarpa*), basswood (*Tilia americana*), green ash (*Fraxinus pennsylvanica*), and eastern cottonwood (*Populus deltoides*; Minnesota Association of Soil and Water Conservation Districts Forestry Committee 1986, Johnson and Larson 1999).

## METHODS

We captured neonatal deer 15 May–15 June from 2002 through 2004 (for model analysis, we included fawns captured and monitored 2001–2002; Brinkman et al. 2004a) using postpartum behavior of reproductive females (Downing and McGinnes 1969, White et al. 1972, Huegel et al. 1985b). Additionally, we captured neonates during 2003 with the aid of VITs (Advanced Telemetry Systems, Isanti, MN) fitted to adults as part of a companion study (Swanson et al. 2008). We fitted captured neonates with expandable breakaway radiocollars (Advanced Telemetry Systems and Telonics Inc., Mesa, AZ). We recorded sex and weight of neonates and determined age (in days) using hoof growth measurement and umbilicus condition (Brinkman et al. 2004b). We calculated parturition date from fawn age at capture and calculated birth mass from age and weight at capture (Nelson and Woolf 1985, Rohm et al. 2007). We located deer daily for the first 8 weeks of life and then 3–4 times/week through 31 August using a truck-mounted null-peak antenna system (Brinkman et al. 2002) and hand-held 4-element Yagi antenna (Advanced Telemetry Systems). When we detected a mortality signal, we immediately (<1 hr) located the collar, conducted field necropsies, and recorded evidence at the site of the mortality to determine cause of death. If we could not determine cause of death in the field, we transported animals to the Animal Disease Research Diagnostic Laboratory (ADRDL) at South Dakota State University for further examination.

Animal handling methods followed guidelines of the American Society of Mammalogists (Gannon et al. 2007) and were approved by the Institutional Animal Care and Use Committee at South Dakota State University (Approval No. 00–A038, 02–A037, and 02–A043).

To determine if habitat characteristics influenced neonate survival, we created 500-m circular buffers around capture locations (Kie et al. 2002, Rohm et al. 2007). Corresponding buffer areas (0.78 km<sup>2</sup>) comprised a land area equivalent to the largest mean 50% summer home range documented for adult females in each of the study areas (Brinkman et al. 2005, Burris 2005, Swanson 2005). To determine habitat characteristics associated with each neonate, we overlaid buffered areas with the 2001 National Land Cover Data set (Homer et al. 2007) and calculated habitat composition (% composition of each buffer) using ArcGIS 9.2 (Esri, Inc., Redlands, CA). We re-classified land cover data into 6 categories, including open water, forested cover, grassland–herbaceous, pasture–hay, cropland, and wetlands. We used FRAGSTATS Version 3.3 to calculate landscape and class-level metrics associated with each buffered area (McGarigal et al. 2002).

We selected the initial set (19 variables) of landscape and class-level metrics important to neonates based upon neonate ecology and metrics significant to survival in previous studies (Vreeland et al. 2004, Rohm et al. 2007). We defined metrics as a) patch density (number of patches/100 ha of the cover type), b) shape index (i.e., average departure of patches from max. compaction), and c) landscape shape index (i.e., standardized measure of the edge for all cover type patches; McGarigal et al. 2002). We tested for colinearity of independent habitat variables using variance inflation (Allison 1999), which resulted in 13 uncorrelated variables that we used to determine the impact of habitat characteristics on neonate survival (Table 1).

To estimate survival to the end of the field season each summer (31 Aug) and determine factors influencing fawn survival, we used known fate models in Program MARK (White and Burnham 1999). We constructed 2 sets of candidate models: 1) model set 1 generated survival estimates and quantified the influence of intrinsic covariates on fawn survival and 2) model set 2 generated survival estimates and quantified the influence of habitat covariates on fawn survival (Rohm et al. 2007). We used the best approximating model from model set 1 as the underlying (constant) structure for all models in model set 2 to account for maximum variation in the data (Burnham and Anderson 2002, Hill et al. 2003, Zablán et al. 2003, Rohm et al. 2007). Intrinsic variables included capture year, sex, site, birth mass, age at death (days), 3-stage age-interval at death (0–2 weeks, 2–8 weeks, and >8 weeks of age; Nelson and Woolf 1987, Rohm et al. 2007), and parturition date relative to peak parturition.

Model set 1 consisted of 13 a priori models (Table 2) constructed from various combinations of intrinsic variables. Model set 2 consisted of 14 a priori models (Table 3; plus best approximating model from model set 1 as the constant survival model) constructed from various combinations of habitat variables. We based a priori model construction on

**Table 1.** Final variables we measured within neonate capture areas used to model the influence of habitat characteristics on neonate white-tailed deer survival in eastern South Dakota and southwestern Minnesota, USA, 2001–2004.

Variable name	No.	Definition
Forested cover	1	Total forested cover (%)
Grassland cover	2	Total grassland cover (%)
Cropland cover	3	Total cropland cover (%)
Wetland cover	4	Total wetland cover (%)
Mean forest patch area	5	Average patch size (ha) for all forest patches
Forest shape index	6	Average departure of forest patches from max. compaction (i.e., square shape)
Mean grassland patch area	7	Average patch size (ha) for all grassland patches
Grassland patch density	8	Density (no./100 ha) of grassland patches
Grassland shape index	9	Average departure of grassland patches from max. compaction (i.e., square shape)
Mean wetland patch area	10	Average patch size (ha) for all wetland patches
Landscape shape index	11	Standardized measure of the amt of edge adjusted for size of the buffered area
Mean patch area	12	Average patch size (ha) for all habitat patches
CV	13	Mean CV of patch size for all habitat patches

**Table 2.** A priori models constructed for model set 1 to determine the influence of intrinsic variables on white-tailed deer neonate survival in eastern South Dakota and southwestern Minnesota, USA, 2001–2004.

Model	$K^a$	Description
$S_{\text{constant}}$	1	Survival was constant
$S_{\text{age}}$	16	Survival varied by age (days) of neonate
$S_{\text{age-interval}}^b$	4	Survival varied by age of neonate in 3 stages
$S_{\text{year}}$	4	Survival varied among the 4 yr
$S_{\text{sex}}$	2	Survival varied between gender of neonates
$S_{\text{site}}$	4	Survival varied between the study sites
$S_{\text{date}}^c$	2	Survival varied between date of birth
$S_{\text{mass}}$	2	Survival varied by birth mass
$S_{\text{age} + \text{mass}}$	17	Survival varied by age and birth mass
$S_{\text{age} + \text{date}}^c$	17	Survival varied by age and parturition date groups
$S_{\text{age} + \text{sex}}$	17	Survival varied by age and sex
$S_{\text{age} + \text{site}}$	19	Survival varied by age and study site
$S_{\text{sex} + \text{year}}$	19	Survival varied by sex and year

<sup>a</sup> Number of parameters.

<sup>b</sup> 3-Stage age-interval: 0–2 weeks, 2–8 weeks, and >8 weeks.

<sup>c</sup> We grouped neonates into peak born and nonpeak born categories.

**Table 3.** A priori models constructed for model set 2 to determine the influence of habitat variables on white-tailed deer neonate survival in eastern South Dakota and southwestern Minnesota, USA, 2001–2004.

Model <sup>a</sup>	Variables <sup>b</sup>	Description
$S_{fc}$	1	% forest cover influences survival
$S_{\text{landscape} + fc}$	1, 11–13	Landscape metrics and % forest cover
$S_{\text{landscape} + \text{grassland}}$	2, 11–13	Landscape metrics and % grassland
$S_{\text{grassland}}$	2	% grassland influences survival
$S_{\text{landscape} + \text{forest metrics}}$	5, 6, 11–13	Landscape metrics and forest metrics
$S_{\text{landscape}}$	11–13	Landscape metrics
$S_{\text{landscape} + \text{grassland metrics}}$	7–9, 11–13	Landscape metrics and grassland metrics
$S_{\text{size}}$	5, 7, 10, 12	Variables pertaining to size influence survival
$S_{\text{landscape} + \text{cropland}}$	3, 11–13	Landscape metrics and % cropland
$S_{\text{age-interval}}$	0	Survival is constant
$S_{\text{wetland}}$	4	% wetland influences survival
$S_{\text{cropland}}$	3	% cropland influences survival

<sup>a</sup> All models have the top model from model set 1  $\{S_{\text{age-interval}}\}$  as the basic structure to which we added habitat covariates.

<sup>b</sup> Variables included in model numbered in Table 1.

variables we considered biologically meaningful to neonate ecology and used Akaike’s Information Criterion corrected for small sample size ( $AIC_c$ ) to select models that best described the data. We compared  $AIC_c$  values to select the most parsimonious model and considered models differing by  $\leq 2 \Delta AIC_c$  from the selected model as potential alternatives (Burnham and Anderson 2002). We used Akaike weights

( $w_i$ ) as an indication of support for each model and used multi-model inference to average parameters across potential models. Models  $\leq 2 \Delta AIC_c$  from the best model were examined to ascertain if they differed by 1 parameter from the best model and had essentially the same maximized log-likelihood. In this case, models with additional parameters were unsupported and noncompetitive and considered

**Table 4.** Top-ranked survival models of neonate white-tailed deer from birth to 31 August in eastern South Dakota and southwestern Minnesota, USA, 2001–2004 from model set 1 (intrinsic covariates) when  $\hat{c}$  (a model term representing overdispersion) was 1.0 (i.e., assumed no dispersion).

Model <sup>a</sup>	AIC <sub>c</sub> <sup>b</sup>	ΔAIC <sub>c</sub> <sup>c</sup>	w <sub>i</sub> <sup>d</sup>	K <sup>e</sup>	Deviance
{S <sub>age-interval</sub> }	211.70	0.00	0.76	4	203.70
{S <sub>age</sub> }	214.82	3.11	0.16	8	198.80
{S <sub>age</sub> + mass}	216.56	4.86	0.07	9	198.54
{S <sub>age</sub> + year}	220.65	8.95	0.01	11	198.62

<sup>a</sup> Composition and description of models are listed in Table 1.

<sup>b</sup> Akaike's Information Criterion corrected for small sample size (Burnham and Anderson 2002).

<sup>c</sup> Difference in AIC<sub>c</sub> relative to min. AIC.

<sup>d</sup> Akaike wt (Burnham and Anderson 2002).

<sup>e</sup> Number of parameters.

potential models only because they contributed an additional parameter (Burnham and Anderson 2002). We evaluated competing models with covariates in which  $\beta$  estimates did not have 95% confidence intervals that included zero (Neter et al. 1996, Barber-Meyer et al. 2008). Because there is no goodness-of-fit test statistic available for known-fate models, we investigated model robustness by artificially inflating  $\hat{c}$  (i.e., a model term representing overdispersion) from 1.0 to 3.0 (i.e., no dispersion to extreme dispersion) to simulate various levels of dispersion reflected in Quasi-AIC<sub>c</sub> (QAIC<sub>c</sub>; Devries et al. 2003, Barber-Meyer et al. 2008).

## RESULTS

During 2002–2004, we captured and radiocollared 78 neonates (6 in 2002, 56 in 2003, 16 in 2004; 41 F, 37 M). Peak parturition occurred on 29 May 2002, 24 May 2003, and 24 May 2004. Age at capture ranged from <1 to 13 days, and 78% of fawns were ≤1 week of age at capture; mean age and weight at capture was 4.3 days (SE = 0.1) and 4.1 kg (SE = 0.4), respectively. Average estimated birth mass was 3.0 kg (SE = 0.1,  $n = 78$ ). Of the 78 neonates, we documented 16 mortalities from time of capture until 31 August: 0 in 2002, 12 in 2003, and 4 in 2004. One neonate died <24 hr postcapture; thus, we censored it from survival analyses. Mean age at death was 24.3 days (SE = 6.8,  $n = 15$ ). Predation was the leading cause of mortality; canid predation accounted for 12 (80.0%) neonate mortalities. We were unable to determine cause of death in the field for the remaining 3 (20.0%) mortalities; however, necropsies conducted at the ADRDL revealed starvation as the likely cause of death.

From model set 1, we considered model {S<sub>age-interval</sub>} as the only model that fit the data (AIC<sub>c</sub> wt = 0.76); remaining models were ≥3.1 ΔAIC<sub>c</sub> units from this model (Table 4). Weight of evidence supporting this model was 4.7 times greater than the next model and >10.9 times greater than remaining models (Table 4). Model {S<sub>age-interval</sub>} had the lowest QAIC<sub>c</sub> when  $\hat{c} = 2.0$  (moderate dispersion; QAIC<sub>c</sub> wt = 0.92) and through  $\hat{c} = 3.0$  (extreme dispersion; QAIC<sub>c</sub> wt = 0.94). The  $\beta$  estimates and 95% confidence intervals for the intercept (9.01, 95% CI = 7.05–10.97), 0–2 weeks (−6.73, SE = 1.04, 95% CI = −8.77 to −4.69), 2–8 weeks (−5.12, SE = 1.08, 95% CI = −7.24 to −3.00), and >8 weeks of age at death (−4.74, SE = 1.16, 95% CI = −7.01 to −2.48) indicated  $\beta \neq 0$  for all intervals; survival was best explained by 3-stage age-intervals. Survival estimates for 0–2 weeks, 2–8 weeks, and >8 weeks were 0.91 (95% CI = 0.90–0.92), 0.98 (95% CI = 0.97–0.98), and 0.99 (95% CI = 0.98–0.99), respectively; overall survival was 0.87 (95% CI = 0.83–0.91). Of 23 mortalities (including those from Brinkman et al. [2004b]) used in covariate models, 14 (60.9%) occurred during the first 2 weeks of life and 6 (26.1%) during 2–8 weeks of life; remaining (13.0%,  $n = 3$ ) mortalities occurred >8 weeks of age.

Using habitat variables, we considered 6 competing models as supporting the data; models were ≤2.0 ΔAIC<sub>c</sub> apart (Table 5). The constant model {S<sub>age-interval</sub>} had the lowest AIC<sub>c</sub> and weight of evidence supporting this model was ≥2.3 times greater than competing models and ≥17.1 times greater for noncompeting models. Model {S<sub>age-interval</sub>} had the lowest QAIC<sub>c</sub> when  $\hat{c} = 2.0$  (moderate dispersion; QAIC<sub>c</sub> wt = 0.32) and through  $\hat{c} = 3.0$  (extreme dispersion;

**Table 5.** Top-ranked survival models of neonate white-tailed deer from birth to 31 August in eastern South Dakota and southwestern Minnesota, USA, 2001–2004 from model set 2 (habitat covariates) when  $\hat{c}$  (a model term representing overdispersion) was 1.0 (i.e., assumed no dispersion).

Model <sup>a</sup>	AIC <sub>c</sub> <sup>b</sup>	ΔAIC <sub>c</sub> <sup>c</sup>	w <sub>i</sub> <sup>d</sup>	K <sup>e</sup>	Deviance
{S <sub>age-interval</sub> }	211.70	0.00	0.32	4	203.70
{S <sub>wetland</sub> }	213.40	1.69	0.14	5	203.39
{S <sub>water</sub> }	213.66	1.96	0.12	5	203.65
{S <sub>cropland</sub> }	213.67	1.96	0.12	5	203.66
{S <sub>grassland</sub> }	213.68	1.97	0.12	5	203.67
{S <sub>forested cover</sub> }	213.71	2.00	0.12	5	203.70

<sup>a</sup> Composition and description of models are listed in Table 2; all models have the base structure of the top model from model set 1 {S<sub>age-interval</sub>}.

<sup>b</sup> Akaike's Information Criterion corrected for small sample size (Burnham and Anderson 2002).

<sup>c</sup> Difference in AIC<sub>c</sub> relative to min. AIC.

<sup>d</sup> Akaike wt (Burnham and Anderson 2002).

<sup>e</sup> Number of parameters.

QAIC<sub>c</sub> wt = 0.33). Although competing models  $\{S_{\text{wetland}}\}$ ,  $\{S_{\text{water}}\}$ ,  $\{S_{\text{cropland}}\}$ ,  $\{S_{\text{grassland}}\}$ , and  $\{S_{\text{forested cover}}\}$  were within  $\leq 2.0 \Delta\text{AIC}_c$ ,  $\beta$  estimates and 95% confidence intervals for these habitat parameters encompassed zero. Furthermore, these models differed from  $\{S_{\text{age-interval}}\}$  by the addition of only 1 parameter and had approximately the same maximized log-likelihood; thus, we excluded these models from consideration.

## DISCUSSION

The neonate survival rate we documented (0.87) was greater than previously reported in southern Illinois (0.70: Nelson and Woolf 1987, 0.59: Rohm et al. 2007), Minnesota (0.49: Kunkel and Mech 1994), Maine (0.40: Long et al. 1998), New Brunswick (0.47: Ballard et al. 1999), and Pennsylvania (0.46: Vreeland et al. 2004). Rohm et al. (2007) attributed elevated predation during their study to increased coyote and bobcat (*Lynx rufus*) abundance. However, in southwestern Lower Michigan, Pusateri-Burroughs et al. (2006) documented similar high survival rates (0.90–0.91), which those authors attributed to alternate food sources for coyotes, landscape composition, and dense ground cover. Although actual coyote density estimates were unknown, we believe the high neonate survival we observed was likely the result of low coyote density in the region based on observations during trapping in southwestern Minnesota (Brinkman 2003).

Similar to other studies documenting cause-specific mortality (Huegel et al. 1985a, Kunkel and Mech 1994, Long et al. 1998, Ballard et al. 1999, Rohm et al. 2007), we determined that canid predation (wolves [*Canis lupus*] were believed absent from South Dakota and southwestern Minnesota) was the major source of neonate mortality. In the Midwest, coyote predation was the main cause of fawn mortality (Huegel et al. 1985a, Benzon 1998), significantly reducing survival (Messier et al. 1986). Canid predation accounted for 50–53% of white-tailed deer neonate mortalities in southern Texas (Cook et al. 1971, Carroll and Brown 1977), 69% in southern Illinois (Nelson and Woolf 1987), 77% in south-central Iowa (Huegel et al. 1985a), 67% in southwest Minnesota (Brinkman et al. 2004a), and 64% in southern Illinois (Rohm et al. 2007). Our results are consistent with previous investigations; coyote predation was the main cause of fawn mortality.

Maternal abandonment because of handling was the proximate cause of starvation-related mortality <3 days postcapture in Pennsylvania (Vreeland et al. 2004). However, starvation or malnutrition as the likely cause of mortality has been documented for fawns up to 37 days postcapture; abandonment even after fawns begin to ruminate might cause starvation or malnutrition related mortality not attributed to handling (Vreeland et al. 2004). Although fawn mortality attributed to starvation occurred 4–7 days postcapture during our study, we have no direct evidence of abandonment due to handling or death of females.

Model selection results with intrinsic covariates indicated that neonate survival was best explained by 3-stage age-interval. Nelson and Woolf (1987), in an area of high forest cover, described the 3-stage age-interval and observed that

neonate mortality was greatest when neonates were becoming more active but were not capable of evading predators (i.e., during the second interval). Neonates can avoid predation when they are <2 weeks of age through cryptic coloration and inactivity, and at >8 weeks of age neonates can escape predators (Nelson and Woolf 1987). However, during our study, survival was lowest during the first 2 weeks of life and highest after neonates reached >8 weeks of age. Our observations support previous hypotheses suggesting the greatest period of vulnerability occurred during the first 2 weeks of life (Rohm et al. 2007).

Model set 2 indicated that variation in neonate survival was not explained by habitat covariates. In southern Illinois, large patches of forested habitat adjacent to several smaller non-forest patches likely provided neonates with cover and concealment (Rohm et al. 2007). However, limited and fragmented forested cover in our study area may have explained why forested habitat did not influence survival in this region. Average forest patch size (0.93 ha) available to neonates was smaller than in southern Illinois, where neonates had larger (by a factor of 4–8 than those in our study) mean forest patch sizes (Rohm et al. 2007).

In southern Illinois, neonate survival was higher in areas with more edge habitat (Rohm et al. 2007). Increased survival in areas with greater edge density may have been an indicator of higher quality habitat; edge habitat likely provided better forage for dams, thereby increasing their condition and neonate survival (Vreeland et al. 2004, Rohm et al. 2007). However, we documented no influence of landscape shape index on neonate survival. During our study, landscape shape indices for neonate survival (3.5) and mortality (3.7) were larger (approx. double) than those documented for survival (2.0) and mortality (1.8) in southern Illinois (Rohm et al. 2007). Furthermore, Rohm et al. (2007) documented that survival areas contained more irregular forest patches than mortality areas in southern Illinois. Irregular boundaries of forested patches may have affected the ability of predators to locate and capture neonates with nonlinear edges being more difficult to search than linear edges (Rohm et al. 2007). However, we documented no influence of irregular forest patches on survival; forest shape indices during our study were similar to those documented in southern Illinois (Rohm et al. 2007).

## MANAGEMENT IMPLICATIONS

We provided the first evaluation of the influence of intrinsic and understudied habitat variables on neonate white-tailed deer survival in the Northern Great Plains. Our study indicated that neonate survival was best explained by neonate age. Variables influencing survival may be area specific; for instance, outside our study area, survival may be dependent on available land cover and juxtaposition of habitats. Knowing when critical periods of fawn survival occur and causes of mortality will aid wildlife managers in identifying the time periods necessary to focus management activities aimed towards increasing neonate survival.

## ACKNOWLEDGMENTS

Funding for this study was provided by Federal Aid to Wildlife Restoration administered by South Dakota Department of Game, Fish and Parks and the Minnesota Department of Natural Resources. Funding also was provided by South Dakota State University, Bend of the River Chapter of Minnesota Deer Hunters Association, Bluffland Whitetails Association, Cottonwood County Game and Fish League, Des Moines Valley Chapter of Minnesota Deer Hunters Association, Minnesota Bowhunters, Inc., Minnesota Deer Hunters Association, Minnesota State Archery Association I, North Country Bowhunters Chapter of Safari Club International, Rum River Chapter of Minnesota Deer Hunters Association, South Metro Chapter of Minnesota Deer Hunters Association, and Whitetail Institute of North America. Additionally, many volunteers and technicians (J. Smith, C. Schultz) greatly contributed to the success of this study. We thank R. Auch and D. Ohlen for reviewing this manuscript. We are grateful to the many landowners who granted us permission to conduct our study on their property. Any mention of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the United States Government.

## LITERATURE CITED

- Allison, P. D. 1999. Logistic regression using SAS: theory and application. SAS Institute Inc., Cary, North Carolina, USA.
- Ballard, W. B., H. A. Whitlaw, S. J. Young, R. A. Jenkins, and G. J. Forbes. 1999. Predation and survival of white-tailed deer fawns in northcentral New Brunswick. *Journal of Wildlife Management* 63:574–579.
- Barber-Meyer, S. H., L. D. Mech, and P. J. White. 2008. Elk calf survival and mortality following wolf restoration to Yellowstone National Park. *Wildlife Monographs* 169.
- Benzon, T. 1998. Mortality and habitat use of white-tailed deer fawns in the central Black Hills, South Dakota 1994–1998. South Dakota Department of Game, Fish and Parks. Pittman–Robertson Game Report W-75-R-34 Pierre, South Dakota, USA.
- Bowman, J. L., and H. A. Jacobson. 1998. An improved vaginal-implant transmitter for locating white-tailed deer birth sites and fawns. *Wildlife Society Bulletin* 26:295–298.
- Brinkman, T. J. 2003. Movement and mortality of white-tailed deer in southwest Minnesota. Thesis, South Dakota State University, Brookings, USA.
- Brinkman, T. J., C. S. DePerno, J. A. Jenks, B. S. Haroldson, and J. D. Erb. 2002. A vehicle-mounted radiotelemetry antenna system design. *Wildlife Society Bulletin* 30:256–258.
- Brinkman, T. J., C. S. DePerno, J. A. Jenks, B. S. Haroldson, and R. G. Osborn. 2005. Movement of female white-tailed deer: effects of climate and intensive row-crop agriculture. *Journal of Wildlife Management* 69:1099–1111.
- Brinkman, T. J., J. A. Jenks, C. S. DePerno, B. S. Haroldson, and R. G. Osborn. 2004a. Survival of white-tailed deer in an intensively farmed region of Minnesota. *Wildlife Society Bulletin* 32:726–731.
- Brinkman, T. J., K. L. Monteith, J. A. Jenks, and C. S. DePerno. 2004b. Predicting neonatal age of white-tailed deer in the Northern Great Plains. *Prairie Naturalist* 36:75–81.
- Brown, A. L., and J. A. Litvaitis. 1995. Habitat features associated with predation of New England cottontails: what scale is appropriate? *Canadian Journal of Zoology* 73:1005–1011.
- Burnham, K. P., and D. R. Anderson. 2002. Model selection and inference: a practical information-theoretic approach. Springer-Verlag, New York, USA.
- Burris, B. M. 2005. Seasonal movements of white-tailed deer in eastern South Dakota and southwestern Minnesota relative to traditional ranges and management unit boundaries. Thesis, South Dakota State University, Brookings, USA.
- Carroll, B. K., and D. L. Brown. 1977. Factors affecting neonatal fawn survival in south-central Texas. *Journal of Wildlife Management* 41: 63–69.
- Carstensen, M., G. D. DelGiudice, and B. A. Sampson. 2003. Using doe behavior and vaginal-implant transmitters to capture neonate white-tailed deer in north-central Minnesota. *Wildlife Society Bulletin* 31:634–641.
- Cook, R. S., M. White, D. O. Tainer, and W. C. Glazener. 1971. Mortality of young white-tailed deer fawns in south Texas. *Journal of Wildlife Management* 35:47–56.
- Dahl, T. E. 1990. Wetlands losses in the United States 1780's to 1980's. U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C., USA.
- DelGiudice, G. D., M. R. Riggs, P. Joly, and W. Pan. 2002. Winter severity, survival, and cause-specific mortality of female white-tailed deer in north-central Minnesota. *Journal of Wildlife Management* 66:698–717.
- DePerno, C. S., J. A. Jenks, S. L. Griffin, and L. A. Rice. 2000. Female survival rates in a declining white-tailed deer population. *Wildlife Society Bulletin* 28:1030–1037.
- Devries, J. H., J. J. Citta, M. S. Lindberg, D. W. Howerter, and M. G. Anderson. 2003. Breeding-season survival of mallard females in the prairie pothole region of Canada. *Journal of Wildlife Management* 67:551–563.
- Dijak, W. D., and F. R. Thompson. 2000. Landscape and edge effects on the distribution of mammalian predators in Missouri. *Journal of Wildlife Management* 64:209–216.
- Downing, R. L., and B. S. McGinnes. 1969. Capturing and marking white-tailed deer fawns. *Journal of Wildlife Management* 33:711–714.
- Dusek, G. L., A. K. Wood, and S. T. Stewart. 1992. Spatial and temporal patterns of mortality among female white-tailed deer. *Journal of Wildlife Management* 56:645–650.
- Gannon, W. L., R. S. Sikes, and the Animal Care and Use Committee of the American Society of Mammalogists. 2007. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy* 88:809–823.
- Gavin, T. A., L. H. Suring, P. A. Vohs, Jr., and E. C. Meslow. 1984. Population characteristics, spatial organization, and natural mortality in the Columbian white-tailed deer. *Wildlife Monographs* 91.
- Gese, E. M., R. L. Ruff, and R. L. Crabtree. 1996. Intrinsic and extrinsic factors influencing coyote predation of small mammals in Yellowstone National Park. *Canadian Journal of Zoology* 74:784–797.
- Hill, M. R., J. R. T. Alisauskas, C. Davison Ankney, and J. O. Leafloor. 2003. Influence of body size and condition on harvest and survival of juvenile Canada geese. *Journal of Wildlife Management* 67:530–541.
- Homer, C., J. Dewitz, J. Fry, M. Coan, N. Hossain, C. Larson, N. Herold, A. McKerrow, J. N. VanDriel, and J. Wickham. 2007. Completion of the 2001 national land cover database for the conterminous United States. *Photogrammetric Engineering and Remote Sensing* 73:337–341.
- Huegel, C. N., R. B. Dahlgren, and H. L. Gladfelter. 1985a. Mortality of white-tailed deer fawns in south-central Iowa. *Journal of Wildlife Management* 49:377–380.
- Huegel, C. N., R. B. Dahlgren, and H. L. Gladfelter. 1985b. Use of doe behavior to capture white-tailed deer fawns. *Wildlife Society Bulletin* 13:287–289.
- Johnson, J. R., and G. E. Larson. 1999. Grassland plants of South Dakota and the Northern Great Plains. South Dakota State University College of Agriculture & Biological Sciences South Dakota Agricultural Experiment Station, Brookings, USA.
- Johnson, R. R., and K. F. Higgins. 1997. Wetland resources of eastern South Dakota. South Dakota State University, Brookings, USA.
- Kie, J. G., R. T. Bowyer, M. C. Nicholson, B. B. Boroski, and E. R. Loft. 2002. Landscape heterogeneity at differing scales: effects on spatial distribution of mule deer. *Ecology* 83:530–544.
- Kunkel, K. E., and L. D. Mech. 1994. Wolf and bear predation on white-tailed deer fawns in northeastern Minnesota. *Canadian Journal of Zoology* 72:1557–1565.
- Kunkel, K. E., and D. H. Pletscher. 2000. Habitat factors affecting vulnerability of moose to predation by wolves in southeastern British Columbia. *Canadian Journal of Zoology* 78:150–157.
- Long, R. A., A. F. O'Connell, Jr., and D. J. Harrison. 1998. Mortality and survival of white-tailed deer *Odocoileus virginianus* fawns on a North Atlantic coastal island. *Wildlife Biology* 4:237–247.

- McGarigal, K., S. A. Cushman, M. C. Neel, and E. Ene. 2002. FRAGSTATS: spatial pattern analysis program for categorical maps. <<http://www.umass.edu/landeco/research/fragstats/fragstats.html>>. Accessed 21 Oct 2009.
- Messier, F., C. Barrette, and J. Huot. 1986. Coyote predation on a white-tailed deer population in southern Quebec. *Canadian Journal of Zoology* 64:1134–1136.
- Minnesota Association of Soil and Water Conservation Districts Forestry Committee. 1986. Minnesota tree handbook. Adventure Publications, Staples, Minnesota, USA.
- Minnesota River Basin Data Center. 2003. Basin fact facts. <<http://mrbdc.mnsu.edu>>. Accessed 17 Feb 2005.
- Nelson, M. E., and L. D. Mech. 1986. Mortality of white-tailed deer in northeastern Minnesota. *Journal of Wildlife Management* 50:691–698.
- Nelson, T. A., and A. Woolf. 1985. Birth size and growth of deer fawns in southern Illinois. *Journal of Wildlife Management* 49:374–377.
- Nelson, T. A., and A. Woolf. 1987. Mortality of white-tailed deer fawns in southern Illinois. *Journal of Wildlife Management* 51:326–329.
- Neter, J., M. H. Kutner, C. J. Nachtsheim, and W. Wasserman. 1996. Applied linear statistical models. Fourth edition. WCB McGraw-Hill, New York, New York, USA.
- Porath, W. R. 1980. Fawn mortality estimates in farmland deer range. Pages 55–63 in R. L. Hinein, S. Nehls, editors. White-tailed deer population management in the north central states. Proceedings of the 1979 Symposium of the North Central Section of the Wildlife Society. Transactions of the 41st Midwest Fish and Wildlife Conference of The Wildlife Society, 10 December 1979, Urbana, Illinois, USA.
- Pusateri Burroughs, J., III, H. Campa, S. R. Winterstein, B. A. Rudolph, and W. E. Moritz. 2006. Cause-specific mortality and survival of white-tailed deer fawns in southwestern lower Michigan. *Journal of Wildlife Management* 70:743–751.
- Rohm, J. H., C. K. Nielsen, and A. Woolf. 2007. Survival of white-tailed deer fawns in southern Illinois. *Journal of Wildlife Management* 71:851–860.
- Schulz, J. H., J. L. Ludwig, and M. Frydendall. 1983. Survival and home range of white-tailed deer fawns in southeastern Minnesota. *Minnesota Research Quarterly* 43:15–23.
- Smith, V. J., C. J. Kopplin, D. M. Fecske, and J. A. Jenks. 2002. The South Dakota GAP Analysis Project. Final Report. Research Work Order No. 65. Department of Wildlife and Fisheries Sciences, South Dakota State University, Brookings, South Dakota, USA.
- Stuart-Smith, A. K., C. J. A. Bradshaw, S. Boutin, D. M. Herbert, and A. B. Rippin. 1997. Woodland caribou relative to landscape patterns in north-eastern Alberta. *Journal of Wildlife Management* 61:622–633.
- Swanson, C. C. 2005. Movement and association of white-tailed deer in southwest Minnesota. Thesis, South Dakota State University, Brookings, USA.
- Swanson, C. C., J. A. Jenks, C. S. DePerno, R. W. Klaver, R. G. Osborn, and J. A. Tardiff. 2008. Does the use of vaginal-implant transmitters impact neonate survival of white-tailed deer? *Wildlife Biology* 13:272–279.
- Thogmartin, W. E., and B. A. Schaeffer. 2000. Landscape attributes associated with mortality events in wild turkeys in Arkansas. *Wildlife Society Bulletin* 28:865–874.
- Vogelmann, J. E., S. M. Howard, L. Yang, C. R. Larson, B. K. Wylie, and N. Van Driel. 2001. Completion of the 1990s National Landcover Data Set for the conterminous United States from Landsat Thematic Mapper data and ancillary data sources. *Photogrammetric Engineering and Remote Sensing* 67:650–662.
- Voigtlander, A. L. 1999. Soil survey of Renville County, Minnesota. United States Department of Agriculture, Natural Resources Conservation Service, Washington, D.C., USA.
- Vreeland, J. K., D. R. Diefenbach, and B. D. Wallingford. 2004. Survival rates, mortality causes, and habitats of Pennsylvania white-tailed deer fawns. *Wildlife Society Bulletin* 32:542–553.
- White, G. C., and K. P. Burnham. 1999. Program MARK: survival estimation from populations of marked animals. *Bird Study* 46(Supplement): 120–138.
- White, M., F. F. Knowlton, and W. C. Glazener. 1972. Effects of dam-newborn fawn behavior on capture and mortality. *Journal of Wildlife Management* 36:897–906.
- Whitlaw, H. A., W. B. Ballard, D. L. Sabine, S. J. Young, R. A. Jenkins, and G. J. Forbes. 1998. Survival and cause-specific mortality rates of adult white-tailed deer in New Brunswick. *Journal of Wildlife Management* 62:1335–1341.
- Zablan, M. A., C. E. Braun, and G. C. White. 2003. Estimation of greater sage-grouse survival in North Park, Colorado. *Journal of Wildlife Management* 67:144–154.

*Associate Editor: David M. Forsyth.*